

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup>:</b> <b>A61K 38/21, 31/70 // (A61K 38/21, 31:70)</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 99/59621</b> <b>(43) International Publication Date:</b> 25 November 1999 (25.11.99)
<b>(21) International Application Number:</b> PCT/US99/07037 <b>(22) International Filing Date:</b> 13 May 1999 (13.05.99) <b>(30) Priority Data:</b> 09/079,566 15 May 1998 (15.05.98) US <b>(71) Applicant:</b> SCHERING CORPORATION [US/US]; 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US). <b>(72) Inventor:</b> ALBRECHT, Janice, K.; 1308 Temple Grove Court, Winter Park, FL 32789 (US). <b>(74) Agents:</b> HOFFMAN, Thomas, D. et al.; Schering-Plough Corporation, Patent Dept. K-6-1 1990, 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GD, GE, HR, HU, ID, IL, IN, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LU, LV, MD, MG, MK, MN, MX, NO, NZ, PL, PT, RO, RU, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UZ, VN, YU, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> COMBINATION THERAPY COMPRISING RIBAVIRIN AND INTERFERON ALPHA IN ANTIVIRAL TREATMENT NAIVE PATIENTS HAVING G CHRONIC HEPATITIS C INFECTION  <b>(57) Abstract</b>  Use of ribavirin and interferon alpha to prepare pharmaceutical compositions for treating antiviral treatment naive patient having chronic hepatitis C infection to eradicate detectable HCV-RNA involving a combination therapy using a therapeutically effective amount of ribavirin and a therapeutically effective amount of interferon alpha for a time period of from 20 up to 50 weeks are disclosed.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

5

COMBINATION THERAPY COMPRISING RIBAVIRIN AND INTERFERON ALPHA IN ANTIVIRAL TREATMENT  
NAIVE PATIENTS HAVING G CHRONIC HEPATITIS C INFECTION

10

## BACKGROUND OF THE INVENTION

The present invention relates to uses of ribavirin and interferon  
alpha to prepare pharmaceutical compositions for treating antiviral  
treatment naive patient having chronic hepatitis C infection to eradicate  
15 detectable HCV-RNA involving a combination therapy using a  
therapeutically effective amount of ribavirin and a therapeutically effective  
amount of interferon-alpha for a time period of from 20 up to 50 weeks.

Chronic infection with hepatitis C virus is an insidious and slow-  
20 progressing disease having a significant impact on the quality of life. It  
can eventually result in cirrhosis of the liver, decompensated liver  
disease and/or hepatocellular carcinoma.

Alpha interferon monotherapy is commonly used to treat chronic  
25 hepatitis C infection. However, this treatment is not always effective and  
sometimes results in intolerable side effects related to the dosage and  
duration of therapy. Ribavirin has been proposed as a monotherapy  
treatment for chronic hepatitis C infection (Thomas et al. AASLD  
Abstracts, Hepatology Vol. 20, NO. 4, Pt 2, Number 440, 1994). However,  
30 this monotherapy treatment has usually been found ineffective.  
Combination therapy of alpha interferon and ribavirin has been  
proposed: (Lai, et al. Symposium to the 9th Biennial Scientific Meeting  
Asian Pacific Association for the Study of the Liver. 1994); Combination  
treatment with interferon alpha-2b and ribavirin for chronic hepatitis C in  
35 patients who have failed to achieve sustained response to interferon  
alone: Swedish experience. *J Hepatology*, 1995;232 (Suppl 2):17-21.

Brouwer JT, Nevens F, Michielsen P, et al.; What options are left when hepatitis C does not respond to interferon? Placebo-controlled Benelux multicentre retreatment trial on ribavirin monotherapy versus combination with interferon. *J Hepatol.* 1994;212 (Suppl 1):S17. Abstract WP2/08.

- 5 Chemello L, Cavalletto L, Bernardinello E, et al. Response to ribavirin, to interferon and to a combination of both in patients with chronic hepatitis C and its relation to HCV genotypes. *J Hepatol.* 1994;212 (Suppl 1):S12. Abstract GS5/29; and The effect of interferon alpha and ribavirin combination therapy in naive patients with chronic hepatitis C, *J. Hepatol.* 10 1995;23(Suppl.2):8-12. Reichard et al. *LANCET* 1998; 351;83-87 disclosed that more chronic hepatitis C patients have a sustained virologic response with a combination of interferon alpha-2b and ribavirin for 24 weeks than with only interferon alpha-2b. Reichard et al. also disclosed that interferon-alpha-2b alone is sufficient to achieve a 15 sustained response in such patients with HCV-RNA serum values above 3 million copies/mL. However, no one has described methods using alpha interferon and ribavirin which eradicate HCV-RNA in any long-term, effective manner for antivirally naive patients having a specific HCV genotype infection.

20

There is a definite need for a method for treating antiviral treatment naive patients having chronic hepatitis C infection with a combination of alpha interferon and ribavirin which eradicates HCV-RNA in any long-term, effective manner.

25

### **SUMMARY OF THE INVENTION**

- The invention involves improved uses of ribavirin and interferon alpha to prepare pharmaceutical compositions for treating antiviral 30 treatment naive patient having chronic hepatitis C infection to eradicate detectable HCV-RNA involving a combination therapy using a therapeutically effective amount of ribavirin and a therapeutically effective amount of interferon-alpha for a time period of from 20 up to 50 weeks.

35

We have discovered that if the antiviral treatment naive patient has HCV genotype 1 infection, or if the antiviral treatment naive patient has an HCV genotype 1 infection, and a viral load of greater than 2 million copies per ml of HCV-RNA by quantitative PCR, then the administration of  
5 the combination therapy is effected for a time period of 40 - 50 weeks, preferably 48 weeks.

We have also discovered that if the antiviral treatment naive patient has is HCV genotype 2 or 3 infection, then the administration of  
10 the combination therapy is effected for a time period of 20 - 30 weeks, preferably 24 weeks.

The present invention involves use of ribavirin for the manufacture of pharmaceutical compositions for treating an antiviral  
15 treatment naive patient having chronic hepatitis C infection to eradicate detectable HCV-RNA by a method comprising administering an effective amount of ribavirin in association with an effective amount of interferon alpha for a time period of 20 up to 50 weeks; and, wherein the antiviral treatment naive patient has a HCV genotype 2 or 3 infection, the  
20 administration of ribavirin in association with interferon alpha is effected for a time period of 20 - 30 weeks; and, wherein the antiviral treatment naive patient has an HCV genotype 1 infection, the administration of ribavirin in association with interferon alpha is effected for a time period of 40 - 50 weeks, preferably 48 weeks.

25

The present invention involves use use of interferon alpha for the manufacture of a pharmaceutical composition for treating an antiviral treatment naive patient having chronic hepatitis C infection to eradicate detectable HCV-RNA by a method comprising administering an effective  
30 amount of alpha interferon alpha in association with an effective amount of ribavirin for a time period of 20 to 50 weeks; and wherein the antiviral

treatment naive patient has a HCV genotype 1 infection, administration of ribavirin in association with interferon alpha is effected for a time period of 40 - 50 weeks, preferably 48 weeks and wherein the antiviral treatment naive patient has a HCV genotype 2 or 3 infection, the administration of  
5 ribavirin in association with interferon alpha is effected for a time period of 20 - 30 weeks, preferably 24 weeks.

The present invention involves the use of both ribavirin and alpha interferon alpha for the manufacture of pharmaceutical compositions for  
10 treating an antiviral treatment naive patient having chronic hepatitis C infection, to eradicate detectable HCV-RNA by a method comprising administering an effective amount of ribavirin in association with an effective amount of alpha-interferon for a time period of 20 up to 50 weeks; and wherein the antiviral treatment naive patient has a HCV  
15 genotype 1 infection administration of ribavirin in association with interferon alpha is effected for a time period of 40 - 50 weeks, preferably 48 weeks; and wherein the antiviral treatment naive patient has a HCV genotype 2 or 3 infection, the administration of ribavirin in association with interferon alpha is effected for a time period of 20 - 30 weeks,  
20 preferably 24 weeks.

The present invention involves use of ribavirin for the manufacture of a pharmaceutical composition for treating an antiviral treatment naive patient having chronic hepatitis C genotype 1 infection to eradicate  
25 detectable HCV-RNA by a method comprising administering an effective amount of ribavirin in association with an effective amount of interferon alpha for a time period of 40 - 50 weeks, preferably 48 weeks.

The present invention also involves use of interferon-alpha for the  
30 manufacture of a pharmaceutical composition for treating an antiviral treatment naive patient having chronic hepatitis C genotype 1 infection to

eradicate detectable HCV-RNA by a method comprising administering an effective amount of interferon alpha in association with an effective amount of ribavirin for a time period of 40 - 50 weeks, preferably 48 weeks.

5

A preferred aspect of the present invention further involves use of both ribavirin and interferon-alpha for the manufacture of pharmaceutical compositions for treating an antiviral treatment naive patient having chronic hepatitis C genotype 1 infection to eradicate detectable HCV-1-  
10 RNA by a method comprising administering an effective amount of ribavirin in association with an effective amount of interferon alpha for a time period of 40 - 50 weeks, preferably 48 weeks.

Thus, another preferred aspect of the present invention further  
15 involves use of both ribavirin and alpha interferon for the manufacture of pharmaceutical compositions for treating an antiviral treatment naive patient having chronic hepatitis C genotype 1 infection, and a viral load of greater than 2 million copies per mL of HCV-1 RNA as measured by HCV-RNA quantitative PCR to eradicate detectable HCV-RNA by a  
20 method comprising administering an effective amount of ribavirin in association with an effective amount of interferon alpha for a time period of 40 - 50 weeks, preferably 48 weeks.

Another preferred aspect of the present invention further involves  
25 use of ribavirin for the manufacture of a pharmaceutical composition for treating an antiviral treatment naive patient having chronic hepatitis C genotype 2 or 3 infection to eradicate detectable HCV-RNA by a method comprising administering an effective amount of ribavirin in association with an effective amount of interferon alpha for a time period of 20 - 30  
30 weeks, preferably 24 weeks.

The present invention also involves use of interferon-alpha for the manufacture of a pharmaceutical composition for treating an antiviral treatment naive patient having chronic hepatitis C genotype 2 or 3 infection to eradicate detectable HCV-RNA by a method comprising administering an effective amount of interferon alpha in association with an effective amount of ribavirin for a time period of 20 - 30 weeks, preferably 24 weeks.

10 The present invention further involves use of both ribavirin and alpha interferon for the manufacture of pharmaceutical compositions for treating an antiviral treatment naive patient having chronic hepatitis C genotype 2 or 3 infection to eradicate detectable HCV-RNA by a method comprising administering an effective amount of ribavirin in association  
15 with an effective amount of interferon alpha for a time period of 20 - 30 weeks, preferably 24 weeks.

The interferon-alpha administered is preferably selected from interferon alpha-2a, interferon alpha-2b, a consensus interferon, a  
20 purified interferon alpha product or a pegylated interferon-alpha-2a or pegylated interferon alpha-2b. More preferably, the interferon-alpha is selected from interferon alpha-2a, interferon alpha-2b, or a purified interferon alpha product and the amount of interferon-alpha administered is from 2 to 10 million IU per week on a weekly, TIW, QOD or daily basis.  
25 In a preferred embodiment, the interferon-alpha administered is interferon-alpha-2b and the amount of interferon-alpha is administered 3 million IU TIW.

Alternatively, the interferon-alpha administered is consensus  
30 interferon and the amount of interferon-alpha administered is from 1 to 20 micrograms per week on a weekly, TIW, QOD or daily basis. In another embodiment, the interferon-alpha administered is a pegylated interferon alpha-2b and the amount of pegylated interferon-alpha-2b administered



is from 0.5 to 2.0 micrograms/kilogram per week on a weekly, TIW, QOD or daily basis. Alternatively, the interferon-alpha administered is a pegylated interferon alpha-2a and the amount of pegylated interferon-alpha-2a administered is from 20 to 250 micrograms/kilogram per week on a weekly, TIW, QOD or daily basis.

The use of interferon alpha-2a or pegylated interferon alpha-2a or interferon alpha-2b or pegylated interferon alpha-2b is preferred.

During the 20 - 30 week and during the 40-50 week time periods, the amount of ribavirin administered is 800 to 1200 mg per day, preferably 800, 1000 or 1200 mg per day, and the amount of interferon alpha-2a or interferon alpha-2b administered is from 2 to 10 million IU per week on a weekly, TIW, QOD or daily basis, more preferably 3 million IU TIW.

#### **DETAILED DESCRIPTION**

Surprisingly, it has been found that, in the case of antiviral treatment naive patients having chronic hepatitis C infection and having HCV genotype 1, or such naive patients having HCV genotype 1 and a viral load of greater than 2 million copies per ml of HCV-RNA by quantitative PCR ("PCR"), combination therapy with a therapeutically effective amount of ribavirin and a therapeutically effective amount of interferon alpha for a time period of at least 20 to 30 weeks results in ten times more patients having no detectable HCV-RNA in their serum at least 24 weeks after termination of therapy compared to by interferon-alpha monotherapy. When the combination therapy is extended to a time period of 40 to 50 weeks, two to three times more patients have no detectable HCV-RNA in their serum at least 24 weeks after termination of combination therapy compared to those treated with the combination therapy for 24 weeks and eight to nine times more patients have no detectable HCV-RNA in their serum at least 24 weeks after termination of combination therapy compared to those treated with interferon-alpha monotherapy for 48 weeks. See Tables 6, 14, 16 & 17, the rate of

sustained virologic response found after using the combination therapy of the present invention depends upon the HCV genotype and the base line viral load as measured by HCV-RNA/qPCR as well as the treatment period of the combination therapy for HCV genotype 1. See Tables 13 & 15. The treatment period of the combination therapy for antiviral treatment naive patients having chronic HCV genotypes 4, 5 and 6 infections is the same as antiviral treatment naive patients having chronic HCV genotype 1. The treatment period of the combination therapy for antiviral treatment naive patients having HCV genotypes 2 and/or 3 is shorter, namely 20 to 30 weeks, preferably 24 weeks. See Tables 7, 13 & 15.

The term "interferon alpha" as used herein means the family of highly homologous species-specific proteins that inhibit viral replication and cellular proliferation and modulate immune response. Typical suitable interferon-alfas include, but are not limited to, recombinant interferon alpha-2b such as Intron-A interferon available from Schering Corporation, Kenilworth, N.J., recombinant interferon alpha-2a such as Roferon interferon available from Hoffmann-La Roche, Nutley, N.J., recombinant interferon alpha-2c such as Berofer alpha 2 interferon available from Boehringer Ingelheim Pharmaceutical, Inc., Ridgefield, CT., interferon alpha-n1, a purified blend of natural alpha interferons such as Sumiferon available from Sumitomo, Japan or as Wellferon interferon alpha-n1 (INS) available from the Glaxo-Wellcome Ltd., London, Great Britain, or a consensus alpha interferon such as those described in U.S. Patent Nos. 4,897,471 and 4,695,623 (especially Examples 7, 8 or 9 thereof) and the specific product available from Amgen, Inc., Newbury Park, CA, or interferon alpha-n3 a mixture of natural alpha interferons made by Interferon Sciences and available from the Purdue Frederick Co., Norwalk, CT., under the Alferon Tradename. The use of interferon alpha-2a or alpha 2b is preferred. Since interferon alpha 2b, among all interferons, has the broadest approval throughout the world for treating chronic hepatitis C infection, it is most preferred. The manufacture of interferon alpha 2b is described in U.S. Patent No. 4,530,901.

The interferon alpha administered is selected from interferon alpha-2a, interferon alpha-2b, a consensus interferon, a purified interferon alpha product or a pegylated interferon-alpha-2a or pegylated interferon alpha-2b.

5

The therapeutically effective amount of interferon alpha-2a, interferon alpha-2b, or a purified interferon alpha administered in association with ribavirin is from 2 to 10 million IU per week on a weekly, TIW, QOD or daily basis.

10

The therapeutically effective amount of interferon-alpha-2b administered is 3 million IU TIW.

When the interferon alpha administered in association with  
15 ribavirin is consensus interferon, the therapeutically effective amount of interferon-alpha administered is from 1 to 20 micrograms per week on a weekly, TIW, QOD or daily basis.

The term "pegylated interferon alpha" as used herein means  
20 polyethylene glycol modified conjugates of interferon alpha, preferably interferon alpha-2a and alpha-2b. The preferred polyethylene-glycol-interferon alpha-2b conjugate is PEG<sub>12000</sub>-interferon alpha-2b. The phrases "12,000 molecular weight polyethylene glycol conjugated interferon alpha" and "PEG<sub>12000</sub>-IFN alpha" as used herein mean  
25 conjugates such as are prepared according to the methods of International Application No. WO 95/13090 and containing urethane linkages between the interferon alpha-2a or -2b amino groups and polyethylene glycol having an average molecular weight of 12000. The pegylated inteferon alpha, PEG<sub>12000</sub> -IFN-alpha-2b is available from  
30 Schering-Plough Research Institute, Kenilworth, NJ.

The preferred PEG<sub>12000</sub>-interferon alpha-2b is prepared by attaching a PEG polymer to the epsilon amino group of a lysine residue

in the interferon alpha-2b molecule. A single PEG<sub>12000</sub> molecule is conjugated to free amino groups on an IFN alpha-2b molecule via a urethane linkage. This conjugate is characterized by the molecular weight of PEG<sub>12000</sub> attached. The PEG<sub>12000</sub>-IFN alpha-2b conjugate is formulated as a lyophilized powder for injection. The objective of conjugation of interferon alpha with PEG is to improve the delivery of the protein by significantly prolonging its plasma half-life, and thereby provide protracted activity of interferon alpha.

Other interferon alpha conjugates can be prepared by coupling an interferon alpha to a water-soluble polymer. A non-limiting list of such polymers include other polyalkylene oxide homopolymers such as polypropylene glycols, polyoxyethylenated polyols, copolymers thereof and block copolymers thereof. As an alternative to polyalkylene oxide-based polymers, effectively non-antigenic materials such as dextran, polyvinylpyrrolidones, polyacrylamides, polyvinyl alcohols, carbohydrate-based polymers and the like can be used. Such interferon alpha-polymer conjugates are described in U.S. Patent No. 4,766,106, U.S. Patent No. 4,917,888, European Patent Application No. 0 236 987, European Patent Application Nos. 0510 356, 0 593 868 and 0 809 996 (pegylated interferon alpha-2a) and International Publication No. WO 95/13090.

Pharmaceutical compositions of pegylated interferon alphasuitable for parenteral administration may be formulated with a suitable buffer, e.g., Tris-HCl, acetate or phosphate such as dibasic sodium phosphate/monobasic sodium phosphate buffer, and pharmaceutically acceptable excipients ( e.g., sucrose), carriers (e.g. human plasma albumin), toxicity agents (e.g. NaCl), preservatives (e.g. thimerosal, cresol or benyl alcohol), and surfactants( e.g. tween or polysorbates) in sterile water for injection. The pegylated interferon alpha-may be stored as lyophilized powders under a refrigeration at 2°-

8°C. The reconstituted aqueous solutions are stable when stored between 2° and 8°C and used within 24 hours of reconstitution. See for example U.S. Patent Nos, 4,492,537; 5,762,923 and 5,766,582. The reconstituted aqueous solutions may also be stored in prefilled, multi-dose syringes such as those useful for delivery of drugs such as insulin. Typical suitable syringes include systems comprising a prefilled vial attached to a pen-type syringe such as the NOVOLET Novo Pen available from Novo Nordisk, as well as prefilled, pen-type syringes which allow easy self-injection by the user. Other syringe systems include a pen-type syringe comprising a glass cartridge containing a diluent and lyophilized pegylated interferon alpha powder in a separate compartment.

When the interferon-alpha administered in association with ribavirin is a pegylated interferon alpha-2b and the amount of interferon-alpha administered is from 0.5 to 2.0 micrograms/kilogram per week on a weekly, TIW, QOD or daily basis.

When the interferon-alpha administered in association with ribavirin is a pegylated interferon alpha-2a and the amount of interferon-alpha administered is from 20 to 250 micrograms/kilogram per week on a weekly, TIW, QOD or daily basis.

Ribavirin, 1-β-D-ribofuranosyl-1H-1,2,4-triazole-3-carboxamide, available from ICN Pharmaceuticals, Inc., Costa Mesa, California, is described in the Merck Index, compound No. 8199, Eleventh Edition. Its manufacture and formulation is described in U.S. Patent No. 4,211,771.

A person suffering from chronic hepatitis C infection may exhibit one or more of the following signs or symptoms:

(a) elevated ALT,

(b) positive test for anti-HCV antibodies,

(c) presence of HCV as demonstrated by a positive test for HCV-RNA,

(d) clinical stigmata of chronic liver disease,

5

(e) hepatocellular damage.

To practice the invention, the combination therapy of interferon alpha and ribavirin are administered to the patient exhibiting one of more  
10 of the above signs or symptoms in amounts sufficient to eliminate or at least alleviate one or more of the signs or symptoms. Interferon alpha formulations, including pegylated interferon alpha formulations, are not effective when administered orally, so the preferred method of administering the interferon alpha or pegylated interferon alpha  
15 formulations is parenterally, preferably by subcutaneous, IV, or IM, injection. The ribavirin is administered to the patient in association with the interferon alpha, that is, the interferon alpha dose is administered during the same period of time that the patient receives doses of ribavirin. Ribavirin may be administered orally in capsule, tablet or liquid form in  
20 association with the parenteral administration of pegylated interferon-alpha. Of course, other types of administration of both medicaments, as they become available are contemplated, such as by nasal spray, transdermally, by suppository, by sustained release dosage form, and by pulmonary inhalation. Any form of administration will work so long as the  
25 proper dosages are delivered without destroying the active ingredient.

The term "antiviral treatment naive patients" in the context of the present invention means that the patients have never been treated with ribavirin or any interferon including, but not limited to an interferon-alpha.

30

The term "no detectable HCV-RNA" in the context of the present invention means that there is less than 100 copies of HCV-RNA per ml of serum of the patient as measured by quantitative, multi-cycle reverse transcriptase PCR methodology. HCV-RNA is preferably measured in the  
35 present invention by the methodology described below. This methodology is referred to herein as HCV-RNA/qPCR.

RNA is extracted from patient serum using a guanidinium thiocyanate- phenol-chloroform mixer followed by ethanol-ammonium acetate precipitation. The precipitated RNA is centrifuged and the  
5 resulting pellet is dried in a Centrivap console (Labconco, Kansas City, Mo.). The dry pellet is then resuspended in 30 microliters of an Rnasin (Promega Corp., Madison, WI), dithiothritol, and diethylpyrocarbonate-treated water mixture. Samples are kept at or below -20°C (preferably below -70°C) until RNA reverse transcription (RT) and PCR.

10

In order to convert the entire RNA sequence into cDNA in the RT reaction, random hexadeoxyribonucleotides (Pharmacia Biotech, Piscataway, NJ) are used as primers for the first strand cDNA synthesis. Two aliquots of 3 microliters of resuspended sample is added to 3  
15 microliters of 100ng/μl random primers and denatured at 70°C, then reverse transcribed at 40°C for one hour using M-MLV reverse transcriptase (USB, Cleveland, OH) in standard buffer containing 5 mM MgCl<sub>2</sub>. The final RT reaction volume is 26 μl. The PCR is started immediately following the reverse transcription.

20

A modified version of the PCR method is performed using heat-stable Taq polymerase to amplify the cDNA. Seventy-five microliters of PCR mix is added to the entire RT reaction volume (26 μl) to a final MgCl<sub>2</sub> concentration of 1.5 mM in a total volume of 101 μl. Each 101 μl sample  
25 is then split into 50.5 μl, and a layer of mineral oil is placed on top to prevent evaporation.

The PCR cycle consists of annealing for 90 sec., extension for 90 sec., and denaturation for 90 sec., at 55°C, 74°C and 94°C, respectively.  
30 Thermocycling samples is submitted to a final 74°C extension for 10 minutes. Four different cycle sets are used. By loading the sample in duplicate, and splitting these samples evenly after RT, there are four tubes from one sample. Each of the four tubes is given a different cycle number, enhancing sensitivity and accuracy in the quantitation process.  
35 The thermocycling efficiency will be assessed by satisfactory amplification of known copy number RNA standards included in each set

of 60 tubes. Two primer sets are used for the amplification, both from the 5' untranslated region of the HCV genome. Both of these primer sets are highly conserved and detect all known subtypes of HCV. Primer set 1: upstream 5' -GTG GTC TGC GGA ACC GGT GAG T-3', downstream 5'-  
5 TGC ACG GTC TAC GAG ACC TC-3' which produced a 190 bp product. Primer set 2: upstream 5'-CTG TGA GGA ACT ACT GTC TTC-3', downstream 5'-CCC TAT CAG GCA GTA CCA CAA-3' which produced a 256 bp product.

10 The amplified cDNA is then electrophorised in 3% agarose gel and transferred to nylon membrane. The target DNA is detected by Southern blotting and immunostaining using a nonradioactive digoxigenin-labeled DNA probe. These procedures are performed using automated instruments for PCR thermocycling, agarose gel  
15 electrophoresis, vacuum-transfer Southern blot, hybridization, and immunostaining. Each membrane contains known copy number serially diluted standards which are used to construct standard curves for quantitative measurement of the specimen bands. Originally standard curves are made from carefully diluted HCV-RNA from transcribed  
20 clones. Radioactive incorporation studies, gel electrophoresis, and OD 260 are performed on the transcripts to determine that they are of the expected length. After the production of the RNA transcripts quantitated clone standards "pooled" standards are generated which better represent the heterogeneous nature of HCV, one would encounter in natural  
25 infection. These pools are made by combining large amounts of serum or plasma from known infected individuals. The serum/plasma pools are calibrated with PCR, against the clone transcripts and then diluted in the known PCR-negative fluids. Finally, the higher copy number samples of the pools are checked against the cDNA Quantiplex nucleic acid  
30 detection system from Chiron Inc. (Emeryville, CA). These "double quantitated" pools are aliquoted and saved at -70°C. Dilutions of 5,000,000, 1,000,000, 500,000, 100,000, 10,000, and 1000 copies/ml are used in each experiment.

35 Each Southern blot membrane is scanned into a computer using an automated scanner/densitometer, at intervals during development to



determine when the standard curve is most linear. The resultant electronic images are then measured for band area and mean band density. All of the reading are standardized to integrated band density and compared to the standard curve to obtain a numerical value of viral copy number for each band.

The term "sustained virologic response" as used in the context of the present invention means that there is no detectable HCV-RNA in the serum of patients treated in accordance with the present invention for at least 24 weeks after the end of the combined therapy treatment. Preferably, the period of sustained virologic response is at least one year - or longer - after the end of treatment.

The following clinical protocols were performed:

15

#### **Study 1:**

##### Overall Design and Plan of the Study

20 This was a prospective, multicenter, randomized, double-blind, parallel-group. The study compared treatment with INTRON® A plus ribavirin to treatment with INTRON® A plus placebo for 24 or 48 weeks in antiviral treatment naive patients with compensated chronic hepatitis C who had no prior treatment with any interferon including but not limited to  
25 alpha interferon (INTRON® A, Roferon®-A, consensus interferon, or Wellferon®) therapy and who also had no prior treatment with ribavirin. Patients who had prior treatment for hepatitis with any other antiviral or immunomodulatory drug within the previous 2 years were also excluded from this study. Eligible patients had chronic hepatitis C confirmed by  
30 positive serum HCV-RNA, liver biopsy, and laboratory tests.

Patients were randomized to treatment with either INTRON® A plus ribavirin or INTRON® A plus placebo. The dose of INTRON® A was 3 million IU SC TIW; the dose of ribavirin was 1000 or 1200 mg PO daily  
35 (based on weight) in two divided doses. Treatment group assignments were made in equal ratios by a Central Randomization Center. The

randomization procedure was designed to attempt to balance the treatment groups, within and across sites, with respect to presence or absence of cirrhosis in the pretreatment liver biopsy, serum HCV-RNA/qPCR level, and HCV genotype.

5

Study treatment was administered for 24 or 48 weeks. The total course of the study was 48 or 72 weeks to determine long-term effect of treatment. Duration of treatment was assigned at the time of randomization.

10

During treatment and posttreatment follow-up, biochemical (ALT), virological (HCV-RNA), and histological (liver biopsy) examinations were used to assess the nature and duration of response to study treatment. The primary efficacy variable was the overall response defined as loss of serum HCV-RNA/qPCR (<100 copies/mL) as measured at 24 weeks following the end of therapy. In addition, a decrease in hepatic inflammation, an improvement in posttreatment liver biopsy as measured by the Knodell Histology Activity index (HAI) and normalization of ALT were also examined as secondary efficacy endpoints. The safety of the study treatments was assessed by monitoring selected laboratory parameters and by also recording and evaluating the occurrence of any adverse events.

15

20

#### Treatment Regimens

25

There were four study treatment regimens :

1. INTRON® A 3 million IU SC TIW plus ribavirin 1000 or 1200 mg/day PO in two divided doses for 24 weeks; or
- 30 2. INTRON® A 3 million IU SC TIW plus placebo matching ribavirin PO in two divided doses for 24 weeks; or
3. INTRON® A 3 million IU SC TIW plus ribavirin 1000 or 1200 mg/day PO in two divided doses for 48 weeks; or
- 35 4. INTRON® A 3 million IU SC TIW plus placebo matching ribavirin PO in two divided doses for 48 weeks.

Study treatments 1 and 2 were administered for 24 weeks; study treatments 3 and 4 were administered for 48 weeks. The standard INTRON® A (interferon alpha-2b, recombinant) regimen for hepatitis C was administered as a fixed dose of 3 million IU TIW. Each patient received instructions regarding the preparation and subcutaneous administration of INTRON® A. Ribavirin was administered twice daily, morning and evening. The dose was determined by the patient's body weight at the Entry visit. Patients weighing  $\leq 75$  kg received 1000 mg daily as two 200 mg capsules in the morning and three 200 mg capsules in the evening. Patients weighing  $> 75$  kg received 1200 mg daily as three 200 mg capsules morning and evening.

The randomization procedure was designed to balance the groups with respect to the following Baseline characteristics:

- pretreatment liver histology (cirrhosis or no cirrhosis);
- serum HCV-RNA/qPCR status (HCV-RNA/qPCR  $\leq 2,000,000$  or HCV-RNA/qPCR  $> 2,000,000$  copies/mL); and
- HCV Genotype (1 or other). Patients with mixed genotypes (which include Type 1) will be classified as Type 1 for purposes of balancing.

#### Efficacy

The primary efficacy objective was comparison of the treatment groups 1 and 2 and 3 and 4 with respect to the sustained virologic response rate defined as loss of (detectable) serum HCV-RNA/qPCR measured at 24 weeks following the end of therapy to an undetectable level or to a level  $< 100$  copies/mL. The following secondary efficacy Endpoints were also examined:

The secondary efficacy Endpoints:

- proportions of patients with normalization of ALT at 24 weeks of follow-up;
- proportions of patients with improvement in biopsy (Categories I + II + III combined scores);
- changes from Baseline in the biopsy scores (Categories I + II + III combined scores);
- response rates at Endpoint of treatment based on HCV-RNA/qPCR;
- proportion of patients with normalization of ALT at Endpoint of treatment.
- response rates at 24 weeks of follow-up based on HCV-RNA/qPCR.

Virology: Entry Status and Change from Entry

Serum HCV-RNA/qPCR testing was performed by a central laboratory. A positive HCV-RNA assay result was required at Baseline; only patients positive for HCV-RNA were eligible to participate. Repeat assays were scheduled at Weeks 4, 12, 24, and if the patient was in the 48 week treatment groups at weeks 36 and 48. All patients had repeat assays scheduled for Follow-up Weeks 12 and 24.

Response was assessed as defined below:

**Responder:** A patient was classified as a responder at a given time point if HCV-RNA/qPCR was negative (<100 copies per mL) at that time point.

**Sustained Responder:** A patient was classified as a sustained responder if the patient was a responder at 24 weeks of follow-up.

Note that patients who do not meet these criteria, including patients who discontinued before the required HCV-RNA/qPCR evaluations were obtained, were classified as non-responders.

**Overall Responder:** Based on both serum HCV-RNA/qPCR and change in liver histology as evaluated by the Knodell HAI Inflammation Score. A patient was classified as an overall responder to treatment if he/she was a sustained responder and his/her Post treatment Knodell HAI inflammation score (sum of categories I+II+III) improved by 2 or more units relative to the Pretreatment score.

#### Liver Histology

Liver biopsy was required within the six months preceding patient enrollment and at Follow-up Week 24 for all patients. Evaluation of the biopsies was performed by a single pathologist using the Knodell Histology Activity Score. The central pathologist was blinded with respect to patient identification, treatment group, and the time the biopsy was obtained relative to treatment (Pre- or Post treatment). Efficacy of study treatments was assessed by comparing the degree of inflammatory activity observed at Baseline with that present at Follow-up Week 24.

#### RESULTS

Nine hundred-twelve patients were enrolled at 42 US centers and randomized to treatment with either INTRON® A plus ribavirin (N=228) or INTRON® A plus placebo (N=23) for 24 weeks or treatment with either INTRON A plus ribavirin ("I+R") (N=228) or INTRON A plus placebo ("I+P") (N=225) for 48 weeks.

Overall, 81% (734/912) of patients completed treatment and 24 weeks of follow-up. Eighty-nine percent (203/228) of patients in the 24 week I+R group, 90% (207/231) of patients in the 24 week I+P group, 70% (159/228) of patients in the 48 week I+R group, and 73% (165/225) of patients in the 48 week I+P group completed the study.

Twenty percent (178/912) of patients discontinued during treatment: 11% (25/228) in the 24 week I+R group, 10% (24/231) in the 24 week I+P group, 30% (69/228) in the 48 week I+R group, and 27% (60/225) in the 48 week I+P group. An adverse event was the most frequent reason a

patient discontinued treatment in all groups (8% [19/228] with 24 weeks of I+R, 9% [20/231] with 24 weeks of I+P, 20% [45/228] with 48 weeks of I+R, and 14% [32/225] with 48 weeks of I+P.

At least 96% of patients who completed treatment and entered follow-up completed the study. Only 2 patients in the I+R 24 week group, 8 patients in the 24 week I+P group, 7 patients in the I+R 48 week group, and 4 patients in the 48 week I+P group discontinued during follow-up.

The patient's weight and their baseline disease characteristics (HCV genotype and initial viral load) for all patients in Study 1 is given in Table 1 below. HCV genotypes were done on the patient serum samples subjected to HCV-RNA/qPCR testing.

**Table 1. Body Weight and Baseline Disease Characteristics for Study 1 Patients**

	I+R <sup>1</sup> 24 Wks (N=228)	I+P <sup>2</sup> 24 Wks (N=231)	I+R <sup>1</sup> 48 Wks (N=228)	I+P <sup>2</sup> 48 wks (N=225)
<b>Weight</b>				
>75 kg	148 (65%)	157 (68%)	133 (58%)	153 (68%)
<75 kg	80 (35%)	74 (32%)	95 (42%)	72 (32%)
<b><u>HCV Genotype<sup>3</sup></u></b>				
1	164 (72%)	167 (72%)	166 (73%)	162 (72%)
2	29 (13%)	38 (17%)	37 (16%)	43 (19%)
3	28 (12%)	24 (10%)	23 (10%)	19 (8%)
4	6 (3%)	2 (0.9%)	1 (0.4%)	1 (0.4%)
5	0	0	1 (0.4%)	0
6	1 (0.4%)	0	0	0
<b><u>HCV-RNA/qPCR</u></b>				
(copies/mL)				
Geometric Mean	3,070,019	2,767,469	2,922,925	2,819,324
≤ 2 million copies/mL	62 (27%)	74 (32%)	76 (33%)	63 (28%)
> 2 million copies/mL	166 (73%)	157 (68%)	152 (67%)	162 (72%)

**FOOTNOTES TO TABLE 1**

1. I + R is Intron A + Ribavirin
2. I + P is Intron A + Placebo
3. Sub-genotypes are classified under their respective genotype.

All discussions of efficacy and safety in this report are based on data for the all-treated groups.

### Efficacy

The objectives of this study were to compare INTRON® A plus  
5 ribavirin with INTRON® A plus placebo with respect to the overall  
response rate and the virologic response rate (based on HCV-RNA  
(qPCR) for 24 and 48 weeks. The primary efficacy variable for the study  
is the overall response rate.

10 The conclusions from this regarding efficacy are as follows:

The combination therapy of INTRON A plus ribavirin administered for  
48 weeks increased by 2 to 3-fold the efficacy of INTRON A monotherapy  
for the treatment of chronic hepatitis C in antiviral treatment naive  
15 patients. Forty-eight weeks of the combination therapy of INTRON A plus  
ribavirin increased the response rate at the End of Treatment and  
decreased the rate of relapse, which resulted in a better sustained  
virologic response rate than for 48 weeks of INTRON A +Placebo. This  
enhancement of efficacy included all aspects of the disease and resulted  
20 in:

- Sustained eradication of detectable **HCV-RNA**;
- Improvement in hepatic inflammation;
- Normalization of ALT;
- Improvement in Knodell HAI inflammation score.

25

Sustained loss of serum HCV-RNA correlated with improvement in  
or resolution of hepatic inflammation. Results demonstrated correlation  
between sustained virologic response, improvement in hepatic  
inflammation, normalization of ALT and improvement in HQL.

30



The End of Follow-up overall response rate is a composite of the loss of serum HCV-RNA(qPCR) and change in liver histology at end of follow-up (24 weeks following the end of treatment). A patient was classified as an overall responder if HCV-RNA(PCR) was negative at the 24 week posttreatment evaluation and the posttreatment Knodell HAI inflammation score (sum of categories I+II+III) had improved if the posttreatment value had decreased by 2 or more units relative to the pretreatment score. The percent of sustained virologic responders by time to first negative HCV-RNA, the End of Follow-up (sustained) virologic response, histologic response, and overall response rates are summarized in **Tables 2, 3, 4 and 5.**

End of Follow-up HCV-RNA Sustained Virologic Response:  
Sustained Loss of HCV-RNA 24 Weeks Following the End of Treatment

15

The proportion of patients with eradication of HCV-RNA in the serum 24 weeks following the End of Treatment was two-threefold greater (41% v. 16%) in the group of patients treated with the combination of INTRON® A plus ribavirin compared to those receiving INTRON® A plus placebo.

20

The increased length of the combination therapy had the greatest effect on relapse rates. At 24 weeks following End of the Treatment, the relapse rates for 48 weeks of the combination therapy and for 48 weeks of Intron A plus placebo were the same (12%). Longer treatment with the combination therapy (48 weeks) and decreased relapse rates resulted in the highest sustained virologic response rates. Sustained virologic response rates were also significantly higher with 48 weeks of the combination therapy compared to 24 weeks of the combination therapy (38% vs 31%, p value = 0.053.)

30

Extending the combination therapy from 24 to 48 weeks substantially increased sustained virologic responses in patients who first became HCV-RNA negative at Weeks 12 and 24. The majority of patients who became sustained virologic responders were HCV-RNA negative by Week 4. As summarized in Table 2, the sustained virologic response was observed for 81% (35/44) of the patients on 24 weeks of combination therapy (I+R) at Week 4 of the 24 week treatment and for 81% (36/45) of the patients on the 48 weeks of combination therapy at Week 4 of the 48 week treatment. Note that substantial portions of these patients on the 24 and 48 week combination therapy who responded for the first time at Week 12 became sustained virologic responders. In 42% of these patients in the 24 week combination therapy group and 63% of these patients in the 48 week combination therapy group, these responses were sustained. Furthermore, 44% of the patients in the 48 week combination therapy treatment group who first became HCV-RNA negative at Week 24 achieved a sustained virologic response. None of the responses that occurred after Week 24 became sustained responders in any treatment group.

The number of first time responders at Weeks 12 and 24 who become sustained virologic responders 24 weeks after end of treatment were greatest for the patients who received 48 weeks of combination therapy. (See Table 2 below.)

**Table 2. Percent of Sustained Virologic Responders by  
Time to First No Detectable HCV-RNA Levels for Study 1**

<u>Time to First No- Detectable HCV- RNA(Weeks)</u>	<u>Intron A + Ribavirin</u>		<u>Intron A + Placebo</u>	
	<u>24 weeks</u>	<u>48 weeks</u>	<u>24 weeks</u>	<u>48 weeks</u>
4	81%(35/43)	80%(36/45)	48%(10/21)	70%(14/20)
12	42%(30/72)	63%(40/63)	9%(3/32)	35%(11/31)
24	46%(5/11)	44%(11/25)	0%(0/22)	22%(4/18)

**Table 3** summarizes the End of Follow-up patient response as indicated by serum HCV-RNA.

**Table 3 End of Follow-up Serum HCV-RNA: Proportion of Patients with Eradication of HCV-RNA at 24 Weeks Following the End of Treatment (EOT)**

Patient Response Status All Treated 95% Confidence Interval	INTRON A + Ribavirin		INTRON A + Placebo		P-Value <sup>1</sup>
	A	B	C	D	
<b><u>At EOT</u></b>	<b>24 weeks</b>	<b>48 weeks</b>	<b>24 weeks</b>	<b>48 weeks</b>	
Negative	121 (53%)	115 (50%)	66 (24%)	54 (24%)	<0.001
Positive	107 (47%)	113 (50%)	165 (71%)	171 (76%)	
<b><u>At End of Follow-up</u></b>					
Sustained Responders	70 (31%)	87 (38%)	13 (6%)	29 (13%)	<0.001
Relapser	54 (24%)	28 (12%)	53 (23%)	26 (12%)	
Non Responder	104 (46%)	113 (50%)	165 (71%)	170 (76%)	

<sup>1</sup>Fisher's Exact Test for End of Treatment: P Value A vs B is 0.639; P Value for A vs C <0.001; P Value for A vs D <0.001.

<sup>2</sup>Logistic Regression for End of Follow-Up Comparisons; P Value A vs B is 0.053; P Value for A vs C <0.001; P Value for A vs D <0.001.

Combination therapy significantly increased the virologic response at the end of treatment compared to INTRON A monotherapy. See Table 3. The p values for comparison of the combination therapy for 48 weeks with INTRON A monotherapy for 48 weeks and the comparisons of 24 weeks of combination therapy with 24 and 48 weeks of INTRON A monotherapy are each <0.001. Extending the combination therapy from 24 to 48 weeks decreased the relapse rate by 50% (24% to 12%) thereby making the 48 week combination therapy more effective than the 24 week combination (p=0.053).

Pre- and Posttreatment biopsies were available for 79% (179/228) and 69% (157/228) of the patients treated with INTRON® A plus ribavirin for 24 and 48 weeks respectively and for 76% (176/231) and 70%

(158/225) of those patients who received INTRON® A plus placebo for 24 and 48 weeks, respectively. **Table 4** summarizes the effect of treatment on hepatic inflammation for patients with both pre- and posttreatment liver biopsy results. As with the sustained loss of HCV-RNA replication, the  
5 proportion of patients with improvement in liver inflammation was significantly greater ( $p < 0.001$ ) in patients receiving combination therapy compared to those receiving INTRON® A monotherapy for 48 weeks.

Eradication of detectable HCV-RNA was highly correlated with  
10 normalization of serum ALT. Two to three-fold more patients were ALT normal with the combination therapy compared to INTRON A monotherapy at the End of Follow-Up. Among patients who had sustained normalized ALT, a higher proportion of 48 week combination therapy patients were sustained virologic responders compared with  
15 patients who received 24 weeks of combination therapy and 24 weeks or 48 weeks of INTRON A monotherapy.

**Table 4 End of Follow-up Liver Histology: Improvement in Liver Histology 24 Weeks Following the End of Treatment Based on the Knodell HAI (I+II+III) Score.**

Patient Status	Number (%) of Patients <sup>b</sup>				P Value <sup>c</sup>
	INTRON A + Ribavirin		INTRON A + Placebo		
	A	B	C	D	
	24 weeks (N=179) <sup>a</sup>	48 weeks (N=157) <sup>a</sup>	24 weeks (N+176) <sup>a</sup>	48 weeks (N=158) <sup>a</sup>	
Improved Biopsy <sup>d</sup>	102 (57%)	96 (61%)	77 (44%)	65 (41%)	<0.001

a N=Number of Patients with paired biopsy samples.

b Patients with both pre- and posttreatment biopsy.

c Fisher's Exact test.

d Change from pretreatment to posttreatment in the Knodell Histological Index (HAI) score (sum of I+II+III) categorized as a decrease of 2 or more from pretreatment.

### Overall Response

5 When the study was designed, it was recognized that because liver biopsy is an invasive procedure that it would be unlikely that posttreatment liver biopsies would be obtained for all patients. Therefore, the protocol and statistical analysis plan specified that the analysis for

10 overall response would be based on data for all treated patients and will be estimated by a maximum likelihood method (MLE) for patients whose overall response status could not be determined, ie, patients with negative HCV-RNA and missing (posttreatment) biopsy evaluations. The protocol also specified that an additional analysis would be performed on

15 patients with both pretreatment and posttreatment biopsy results (ie, patients with complete data). The overall response is a composite of sustained loss of detectable HCV-RNA and improvement in liver histology at End of Follow-Up. Overall response is summarized in **Table 5** based on the following analyses:

- 20
- maximum likelihood estimate (MLE);

- patients with complete data (results for both pre- and posttreatment biopsy);
- patients with missing data (either or both HCV/biopsy) treated as failures.

5

**Table 5** Overall Response Rate.

Data Analyzed	INTRON A + Ribavirin		INTRON A + Placebo		p value <sup>b</sup> B vs D
	A 24 weeks	B 48 weeks	C 24 weeks	D 48 weeks	
Maximum likelihood estimate <sup>a</sup>	26%	35%	5%	9%	<0.001
Patients with complete data <sup>c</sup>	29% (52/179)	41% (64/157)	5% (8/170)	11% (18/158)	<0.001
Treat missing as failures <sup>d</sup>	23% (52/228)	28% (64/228)	28% (64/228)	8% (18/225)	<0.001

a MLE based on logistic regression. Logistic regression analysis of patients with complete biopsy data. The p value for A vs D is <0.001 and for A vs B is 0.096.

b Fisher's exact test.

c Complete data = pre- and posttreatment biopsy results. (Logistic regression analysis.)

d Patients who had either virology or biopsy data missing or both were counted as non-responders.

As would be anticipated from individual results for effect of treatment on eradication of HCV-RNA at end of follow-up and improvement in hepatic inflammation, the overall response rate in the INTRON® A plus ribavirin 48 week treatment group was significantly greater (<0.001), than that observed in the INTRON® A plus placebo 48 week treatment group. There was a statistically significant improvement in the overall response with the 48 week combination therapy treatment compared to 48 weeks of INTRON A monotherapy treatment as measured by MLE and complete biopsy. The overall response rates for 24 and 48 weeks of combination therapy treatment was significantly higher than with 24 and 48 weeks of INTRON A monotherapy, respectively.

Logistic regression analysis was done on all baseline demographic variables and disease characteristics. The only baseline statistically significant patient and disease characteristics predictive of  
5 End of Follow-up sustained response were genotype other than 1 and viral load  $\leq 2$  million.

For number of viral copies ( $\leq 2$  million,  $>2$  million), the difference was statistically significant in favor of higher response rates in patients  
10 with  $\leq 2$  million copies (**Table 6** ).

When genotype and baseline virus load are combined, a hierarchy of response is observed. Those patients with genotype other than 1 and baseline virus load  $\leq 2$  million copies who received 24 and 48 weeks of  
15 combination therapy had the best End of Follow-up response; the sustained virologic response for those patients with genotype 1 and  $>2$  million copies who had 48 weeks of the INTRON A plus ribavirin combination therapy was two times better than those same type patients who had the combination therapy for only 24 weeks. (See Table 6)



**Table 6 Disease Characteristics vs Sustained Response: All-Treated Patients.**

Disease Characteristic <sup>a</sup>	Number (%) of Patients			
	INTRON A + Ribavirin		INTRON A + Placebo	
	24 weeks	48 weeks	24 weeks	48 weeks
<u>HCV-RNA/viral load</u>				
≤ 2 million copies/ml	42% (26/62)	43% (33/76)	9% (7/74)	29% (18/63)
> 2 million copies/mL	27% (44/166)	36% (54/152)	4% (6/157)	7% (11/162)
<u>HCV Genotype</u>				
1	16 % (26/164)	28% (46/166)	2% (3/167)	7% (11/162)
Other Genotypes	69% (44/64)	66% (41/62)	16% (10/64)	29% (18/63)
<u>Genotype/Baseline</u>				
<u>HCV-RNA</u>				
Other Genotypes	88% (14/16)	71 % (15/21)	25% (5/20)	50% (10/20)
≤2 million copies/ml				
Other Genotypes	63% (30/48)	63% (26/41)	11% (5/44)	19% (8/43)
>2 million copies/ml				
Genotype 1, ≤2 million	26% (12/46)	33 % (18/55)	4 % (2/54)	19 % (8/43)
copies/ml				
Genotype 1, >2 million	12% (14/118)	25 % (28/111)	1% (1/113)	3% (3/119)
copies/ml				

a At entry, patients were stratified by number of HCV viral copies (≤2 million, >2 million), genotype (1 or other), and cirrhosis (present or absent) as measured by HCV-RNA/of PCR.

b

5

**Table 7. Sustained Virologic Response Rates (%) by HCV-Genotype**

Genotype	INTRON A + Ribavirin		INTRON A + Placebo	
	<u>24 weeks</u>	<u>48 weeks</u>	<u>24 weeks</u>	<u>48 weeks</u>
1	16% (27/165)	28% (46/166)	21% (4/168)	7% (11/162)
2	83% (25/30)	68% (25/37)	18% (7/38)	35% (15/43)
3	57% (16/28)	65% (15/23)	8% (2/24)	16% (3/19)
4-6	40% (2/5)	50% (1/2)	0	0

10

**Table 7** illustrates that the sustained virologic response rates for patients of each genotype with 48 weeks of the combination therapy were greater than those treated with Introna plus placebo for 24 and 48 weeks. With the exception of HCV-genotype 2 patients, extending the

duration of the combination therapy increased the proportion of patients with sustained virologic responses. But see Table 15 for combined virologic responses for Studies 1 & 2). Logistic regression analysis of sustained virologic response demonstrated that, in addition to treatment group, HCV genotype other than 1 and  $\leq 2$  million copies of baseline HCV virus/mL were significant predictors of sustained virologic response (p value  $\leq 0.0111$ ). Most notably, treatment with the combination therapy for 48 weeks improved the sustained virologic response rates for patients expected to have the lowest response rates, namely, those patients with HCV-genotype 1 and greater than 2 million copies of HCV virus/mL. These patients had sustained virologic response rates which were two fold greater than those who had 24 weeks of treatment with the combination therapy. Significantly the sustained virologic response for HCV-genotype 1 patients who received 48 weeks of combination therapy was 1.75 times greater than that for those who received 24 weeks of combination therapy.

### Study 2:

By basically the same methodology as described above in Study 1, Study 2 was also conducted in 43 international sites (832 patients) using the following three treatment regimens:

The results are summarized below:

1. INTRON® A 3 million IU SC TIW plus ribavirin 1000 or 1200 mg/day  
PO in two divided doses for 24 weeks ; or
2. INTRON® A 3 million IU SC TIW plus ribavirin 1000 or 1200 mg/day  
PO in two divided doses for 48 weeks ; or
3. INTRON® A 3 million IU SC TIW plus placebo matching ribavirin  
PO in two divided doses for 48 weeks

### Efficacy

- 5       The primary efficacy objective is sustained virologic response as defined by the loss of detectable serum HCV-RNA(qPCR) measured at End of Follow-up (24 weeks following the end of treatment). A patient was classified as an overall responder if HCV-RNA(PCR) was negative at the 24 week posttreatment evaluation and the posttreatment Knodell HAI inflammation score (sum of categories I+II+III) had improved (decreased)
- 10       by 2 or more units relative to the pretreatment score. The percent of sustained virologic responders by time to first negative HCV-RNA, the End of Follow-up virologic response, histologic response, and overall response rates are summarized in **Tables 9, 10, 11 and 12.**
- 15       Patient body weight and their baseline disease characteristics (HCV genotype and initial viral load) for all patients in Study 2 is given in **Table 8** below.

**Table 8. Body Weight and Baseline Disease Characteristics for Study 2 Patients**

5

	<b>INTRON A + Ribavirin</b>		<b>INTRON A + Placebo</b>
<b>Body Weight</b>	<b><u>24 weeks</u></b>	<b><u>48 weeks</u></b>	<b><u>48 weeks</u></b>
>75 kg	116 (42%)	129 (47%)	122 (44%)
≤75 kg	161 (58%)	148 (53%)	156 (56%)
<b>HCV Genotype</b>			
1	161 (58%)	159 (57%)	168 (60%)
2	27 (10%)	23 (8%)	21 (8%)
3	73 (26%)	74 (27%)	79 (28%)
4	12 (4%)	16 (6%)	9 (3%)
5	1 (0.4%)	5 (2%)	1 (0.4%)
6	3 (1%)	0	0
<b>HCV-RNA/of PCR (copies/ml)</b>			
mean geometric	2,229,797	2,064,959	2,351,824
≤2 million copies/mL	108 (39%)	115 (42%)	95 (34%)
>2 million copies/mL	169 (61%)	162 (59%)	183 (66%)

**Table 9. Percent of Sustained Virologic Responders by Time to First No Detectable HCV-RNA Levels for Study 2**

<u>Time to First No-Detectable HCV-RNA (Weeks)</u>	<u>Intron A + Ribavirin</u>		<u>Intron A+ Placebo</u>
	<u>24 weeks</u>	<u>48 weeks</u>	<u>48 weeks</u>
4	84%(57/68)	83%(58/70)	72%(33/46)
12	47%(36/77)	69%(51/74)	34%(18/53)
24	12%(3/26)	45%(9/20)	10%(2/21)

5

As summarized in Table 9, the majority of patients who became sustained virologic responders had negative HCV-RNA levels by Week 4 of treatment. However, substantial proportions of patients in the 24 and 48 week combination therapy treatment groups who were HCV-RNA positive at Week 4 responded for the first time at Week 12; 47% of those in the 24 Week treatment group and 69% of those in the 48 Week treatment group became sustained responders. Importantly, 45% (9/20) of the patients in the 48 Week combination therapy treatment group who first became negative at Week 24 became sustained virologic responders. None of the responses that occurred after Week 24 became sustained in any of the three treatment groups.

20

**End of Follow-up HCV-RNA Response: Sustained Loss of HCV-RNA 24 Weeks Following the End of Treatment**

The proportion of patients with eradication of HCV-RNA in the serum 24 weeks following the end of the combination therapy treatment was significantly greater in patients treated with the combination therapy of INTRON® A plus ribavirin compared to those receiving INTRON® A

monotherapy. **Table 10** summarizes the End of Follow-up patient response as indicated by serum HCV-RNA.

**Table 10. End of Follow-up Serum HCV-RNA: Proportion of Patients with Eradication of HCV-RNA at 24 Weeks Following the End of Treatment. (EOT) and End of Follow-Up (EOFU).**

	Number (%) of Patients			p value B vs C <sup>1</sup>
	INTRON A + Ribavirin A 24 weeks (N=277)	B 48 weeks (N=277)	INTRON A + Placebo C 48 weeks (N=278)	
EOT <sup>2</sup>				
Negative	57% (157)	52% (145)	33% (93)	<0.001
Positive	42% (120)	42% (132)	65% (185)	
EOFU <sup>3</sup>				
Sustained	35% (96)	43% (118)	19% (53)	<0.001
Relapse	23% (23)	10% (27)	15% (41)	
Non Responders	42% (42)	48% (132)	66% (184)	

<sup>1</sup>Fisher's Exact Test for EOT comparisons; logistic regression for EOFU.

<sup>2</sup>EOT = End of Treatment. P value for A vs C = <0.001 and for A vs B is 0.348.

<sup>3</sup>EOFU = End of Follow-Up. P value for A vs C = <0.001 and for A vs B is 0.055.

**Table 10** illustrates that (1) the 24 and 48 week combination therapies significantly increased the sustained virologic response at the end of treatment compared to that for the Intron A monotherapy (p values are both <0.001) and (2) the increasing the length of the Combination Therapy from 24 to 48 weeks had the greatest effect on relapse rates (10% for 48 weeks vs 23% for 24 weeks, p value is 0.055.)

10 End of Follow-up Liver Histology: Improvement in Liver Histology  
24 Weeks Following the End of Treatment Based on Knodell Histological  
Activity Index (HAI) Scores (I+II+III)

Pre- and Posttreatment biopsies were available for 74% (204/277) and 60% (167/277) of the patients treated with INTRON® A plus ribavirin for 24 and 48 weeks, respectively, and for 69% (191/278) of those patients who received INTRON® A plus placebo. **Table 11** summarizes the effect of treatment on hepatic inflammation for patients with both pre- and posttreatment liver biopsy results. As with the sustained loss of HCV-RNA replication, the proportion of patients with improvement in liver inflammation was significantly greater (p<0.001) in patients receiving combination therapy for 48 weeks compared to those receiving INTRON® A monotherapy for 48 weeks. Extending the combinatin therapy from 24 to 48 weeks also significantly increased the proportion of patients who had improvement in hepatic inflammation (p value=0.046).

**Table 11. End of Follow-up Liver Histology: Improvement in Liver Histology 24 Weeks Following the End of Treatment Based on the Knodell HAI (I+II+III) Score.**

Patient Status	INTRON A + Ribavirin		INTRON A + Placebo	p value <sup>c</sup> B vs C
	A 24 weeks (N=204) <sup>a</sup>	B 48 weeks (N=167) <sup>a</sup>	C 48 weeks (N=191) <sup>a</sup>	
Improved Biopsy <sup>d</sup>	53% (107)	63% (105)	39% (74)	<0.001

a Number of patients with paired biopsies.

b Patients with both pre- and posttreatment biopsy.

c Fisher's Exact test. P value for A vs C = 0.007 and A for A vs B is 0.046.

d Change from pretreatment to posttreatment in the Knodell Histological Index (HAI) score (sum of I+II+III) categorized as a decrease of 2 or more from pretreatment.

### Overall Response

5

The overall response is summarized in **Table 12** based on the following analyses:

- maximum likelihood estimate (MLE);
- patients with complete data (results for both pre- and posttreatment biopsy);
- 10 • patients with missing data (either or both HCV-RNA/biopsy) treated as failures.



**Table 12. Overall Response Rate.**

Data Analyzed	INTRON A + Ribavirin		INTRON A + Placebo
	<u>A</u> <u>24 weeks</u>	<u>B</u> <u>48 weeks</u>	<u>C</u> <u>48 weeks</u>
ML Estimate	28%	37%	17%
Patients with Complete Biopsy Data <sup>c</sup>	30% (62/204)	41% (68/167)	17% (32/191)
Treat Missing as Failures <sup>d</sup>	22% (62/277)	24% (68/277)	12% (32/278)

a MIE based on logistic regression. P value for B vs C = <0.001 and for A vs C = <0.002 and for A vs B is 0.043.

b Fisher's Exact test.

c Complete data = pre- and posttreatment biopsy results. P value for B vs C is <0.001, for A vs C is <0.005 and for A vs B is 0.032.

d Patients who had either virology or biopsy data missing or both were counted as failures..

5

As would be anticipated from individual results for effect of treatment on eradication of HCV-RNA at End of Follow-up and improvement in hepatic inflammation, the overall response rate in the INTRON A plus ribavirin group is significantly greater with a two fold improvement over that observed with INTRON A plus placebo groups for all methods of evaluation.

Logistic regression analysis was done on all baseline demographic variables and disease characteristics. The only baseline statistically significant characteristic predictive of End of Follow-up sustained response was genotype other than 1.

**Table 13. Sustained Virologic Response Rates by HCV Genotype for Study 2 Patients**

<u>Genotype</u>	<u>Intron A + Ribavirin</u>		<u>Intron A + Placebo</u>
	<u>24 Weeks</u>	<u>48 Weeks</u>	<u>48 Week</u>
1	18%(29/161)	30% (48/159)	11% (19/168)
2	59% (16/27)	74% (17/23)	35% (8/23)
3	66% (48/73)	61% (45/74)	32% (25/79)
4-6	19% (3/16)	38% (8/21)	12% (1/8)

The combination therapy provided higher sustained virologic response rate for all genotype compared to Intron A plus Placebo.

- 5 Extending the duration of the combination therapy to 48 weeks increased the proportion of sustained virologic response for all genotypes except type 3. (See Table 13 as well as Table 15 for the combined vivologic response for studies 1 and 2)

- 10 For number of viral copies ( $\leq 2$  million,  $> 2$  million), there was a numerical difference in favor of higher sustained virologic response rates in patients with  $\leq 2$  million copies (**Table 14**). When genotype and baseline virus load are combined, a hierarchy of response is observed. Those patients with genotype other than 1 and baseline virus load  $\leq 2$
- 15 million copies had the best End of Follow-Up response; those patients with genotype 1 and  $> 2$  million copies who received the longer 48 week treatment with combination therapy had the most significant improvement in substantial virologic response of all the groups.

**Table 14 Disease Characteristics vs Sustained Virologic Response: All-Treated Patients in Study No. 2**

Disease Characteristic	Number (%) of Patients		
	INTRON® A	plus ribavirin	INTRON® A plus Placebo
	<u>24 weeks</u>	<u>48 weeks</u>	<u>48 weeks</u>
<u>HCV-RNA/qPCR</u>			
≤ 2 million copies/mL	44% (48/108)	47% (54/115)	31% (15/49)
> 2 million copies/mL	28% (48/109)	40% (64/102)	13% (24/183)
<u>HCV Genotype</u>			
1	18% (29/161)	30% (48/159)	11% (19/168)
Other Genotype	58% (67/116)	59% (70/118)	31% (34/110)
<u>Genotype/Baseline HCV-RNA/qPCR</u>			
Other genotype /≤ 2 million copies/mL	53% (28/53)	62% (34/55)	31% (15/49)
Other genotype /> 2 million copies/mL	62% (39/63)	57% (36/63)	31% (19/61)
Genotype1&≤ 2 million copies/mL	36% (20/55)	33% (20/60)	30% (14/46)
Genotype1&> 2 million copies/mL	8% (9/106)	28% (28/99)	4% (5/122)

Table 14 illustrates that extending the combination therapy to 48 weeks generally improved the sustained virologic response rates. The highest sustained virologic response rates were observed with patients who received the combination therapy for 48 weeks with genotypes other than 1 and initial HCV levels of ≤2 million copies/mL. Importantly, for patients with genotype 1 and HCV levels >2 million, sustained virologic response rates with 48 weeks of combination therapy were more than 3 times higher than rates with only 24 week of the combination.

### **Conclusions on Efficacy for Combined Results for All**

#### **Patients in Studies 1 + 2**

5           Combination therapy was significantly more efficacious than  
INTRON®A monotherapy for initial treatment of chronic hepatitis C.  
Sustained virologic response rates in these antiviral treatment naive  
patients were almost three times higher with 48 weeks of combination  
therapy compared with 48 weeks of Intron A monotherapy and  
10 significantly higher with 48 weeks compared to 24 weeks of the  
combination therapy. The improvement in sustained response rates  
could be accounted for by two treatment effects: higher response rates at  
the end of treatment and decreased relapse rates. The net result of both  
of these effects was the highest sustained response rate with the 48  
15 weeks of combination therapy compared to 48 weeks of monotherapy or  
a shorter regimen with the combination. This enhancement of efficacy  
with 48 weeks of combination therapy also included other measures of  
response such as biochemical (ALT) and histological endpoints.

20           In fact, sustained loss of serum HCV-RNA was highly correlated  
with other clinical endpoints - normalization of ALT and improvement in  
or resolution of hepatic inflammation. Loss of detectable HCV-RNA at the  
end of follow-up was associated with normalization of ALT in all treatment  
groups but was somewhat higher with combination therapy. The majority  
25 of combination therapy patients who were ALT normal were also HCV-  
RNA negative (83-87%).

Increased length of therapy had the greatest effect on relapse  
rates. At End of Follow Up, the relapse rates for both the 48 week  
30 combination and monotherapy treatment groups were lower than the

relapse rate of the 24 week combination therapy treatment group. The combination of the high end of treatment response resulting from 24 and 48 weeks of the combination therapy compared to 48 weeks of Intron A monotherapy and decreased relapse rates resulted in the highest sustained response rates. Sustained response rates were twice as high with 48 weeks of combination therapy compared with Intron A monotherapy(48) ( $p<0.001$ ). Sustained virologic response rates were also higher with 48 weeks of combination therapy compared to only 24 weeks of the ( $p=0.008$ ).

10

The benefit of combination therapy was maintained irrespective of standard predictors of response to INTRON®A monotherapy and 24-week combination therapy in relapsed patients. At entry to these trials, patients were stratified by the following disease characteristics: HCV genotype (type 1 or other genotypes); extent of HCV virus level (number of virus copies in the serum  $\leq 2$  million/mL or  $>2$  million/mL); and cirrhosis (present or absent). Logistic regression analysis of sustained virologic response demonstrated that, in addition to treatment group, only HCV genotype was a significant predictor of sustained virologic response. Neither pretreatment HCV virus level nor presence of cirrhosis appeared to influence the ability of previously untreated patients to achieve sustained virologic response to the combination therapy.

With 48 weeks of combination therapy, sustained response rates were consistently higher than those for 48 weeks of Intron A monotherapy regardless of genotype and were generally higher than 24 weeks of combination therapy. As a group, patients infected with genotype 1 have been shown to be less responsive to INTRON®A monotherapy than patients infected with other genotypes. Despite this, sustained virologic response rates for genotype 1 were approximately 3 times higher with 48

weeks of combination therapy compared to 48 weeks of Intron A monotherapy and almost twice as high with 48 weeks compared to 24 weeks of combination therapy. The combination therapy consistently produced higher sustained virologic response rates in patients infected with other genotypes. Response rates with both 24 and 48 weeks of combination therapy were higher than with 48 weeks of Intron A monotherapy and in all genotypes except types 2 & 3 which genotypes had the same response rate i.e., about 64% at the end of 24 weeks as well as after 48 weeks of combination therapy), extending the duration of the combination therapy to 48 weeks increased the proportion of patients with sustained virologic responses. (See Table 15)

The combination therapy was also more effective than 48 weeks of Intron A monotherapy in producing sustained virologic responses regardless of virus level at Baseline. Forty-eight weeks of combination therapy consistently produced higher sustained virologic response rates at every level of virus infection than 48 weeks of Intron A monotherapy. Sustained response rates were similar for most virus levels in both the 24 and 48 week combination therapy treatment groups, except with the highest virus levels ( $5 < 6$  and  $\geq 6 \times 10^6$  copies/mL) where sustained response rates with the 48 week combination therapy groups were approximately twice as high as the 24 week combination therapy groups.

As noted previously, patients with genotypes other than type 1 had higher sustained virologic response rates than those with type 1 and a virus level  $\leq 2$  million/mL was associated with a better response rate than  $> 2$  million/mL. It was notable that extending the combination therapy treatment to 48 weeks improved the sustained response rates for patients expected to have the lowest response rates, namely, those with genotype 1 and  $> 2$  million copies of HCV-RNA/mL. Extending the combinatin

therapy to 48 weeks in this group of patients produced sustained virologic response rates that were 4 times higher than those with only 24 weeks of combination therapy.

- 5           Other demographic/disease history characteristics had little effect on outcome with combination therapy. In contrast, considerably lower sustained response rates were noted with 48 weeks of Intron A monotherapy in patients older than 55 years, >75 kg or who were infected through transfusion. All had sustained response rates in the  
10   range of 10-12%.

- Availability of paired biopsies was high compared with similar types of studies in chronic hepatitis C patients. As anticipated, pre- and/or posttreatment liver biopsies were unavailable for a proportion of  
15   patients for a variety of reasons. However, paired biopsies were obtained from 71% of patients. Improvement was noted in a significantly higher proportion of 48 week combinatin therapy patients compared with the 48 week Intron A monotherapy patients at End of Follow Up ( $p<0.001$ ). Twenty-four weeks of combinatin therapy also was  
20   significantly more efficacious than 48 weeks of Intron A monotherapy in improving hepatic inflammation. As noted previously, the correlation with virologic response was maintained whether biopsies were assessed by improvement or mean change from baseline in necroinflammatory score; 64-69% of patients had improvement in hepatic inflammation with mean  
25   changes from baseline of -3.8 to -5.0. The most substantial mean change was in the 48 week Intron A monotherapy patients.

- As anticipated, sustained virologic responders in all treatment groups experienced greater improvement in liver biopsy inflammation  
30   scores than patients who remained HCV-RNA positive, and although the extent of improvement was similar in all groups the proportion of virologic

responders with histologic improvement was at least twice as high with both the 24 and 48 weeks of combination therapy than with 48 weeks of Intron A monotherapy. Extending the combination therapy resulted in higher mean improvements in hepatic inflammation. It is also interesting  
5 to note that patients who relapsed from the 48 week combination therapy had considerable mean improvement in inflammation. The combined results are summarized in Tables 15 to 20.



**Table 15. Combined Virologic Response for Studies 1 and****2**

**Efficacy: Loss of Delectable HCV-RNA at End of Treatment (EOT and Sustained Virologic Response At End of Follow Up (EOFU) and Percent Patients Responding**

No. of Patients <sup>1</sup>	<u>Intron A + Ribavirin</u>		<u>Intron A + Placebo</u>	
	24 weeks	48 weeks	24 weeks	48 weeks
	N=505 A	N=505 B	N=231 C	N=503 D
EOT <sup>2</sup>	53%	50%	29%	24%
EOFU <sup>3</sup>	33%±4	41%±5	6%	16%
HCV Genotype	(No. of Patients)	(No. of Patients)		
Type 1	17%±4 (56/326)	29%±5 (94/325)	2%	9%
2	73%±12 <sup>4</sup> (41/57)	70%±12 <sup>4</sup> (42/60)	18%	35%
3	63%±9 <sup>4</sup> (64/101)	62%±10 <sup>4</sup> (60/97)	8%	29%
Types 4/5/6	24%±10 (5/21)	39%±20 (9/23)	0%	11%

<sup>1</sup> Patient characteristics: (a) 66% male; 34% female; (b) mean age: 42.7 yrs; (c) Mean pretreatment HCV-RNA levels <2 million copies/ml: 34%, and >2 million copies/ml: 66%; Super Quant, NGI; (d) HCV genotype 1=66%; genotype 2=13%, non 1-3=3%.

<sup>2</sup> EOT is End of Treatment: P Value B vs D= <0.001; P Value A vs D= <0.001.

EOFU is End of Follow-Up - 24 weeks posttreatment; P Value A vs B= 0.257; P Value B vs D= <0.001; P Value A vs D= <0.001

<sup>3</sup> Sustained virologic response at End of Follow-Up; P Value A vs B = 0.008

<sup>4</sup> The sustained virologic response for HCV genotypes 2 and 3 at the end of 24 weeks was about 64.5% and at the end of 48 weeks was about 65%.

**Table 16. Percentage of Sustained Virologic Response for All HCV-Genotype 1 Patients**

5

<u>Patient</u>	<u>Intron A + Ribavirin</u>		<u>Intron A + Placebo</u>	
	<u>24 weeks</u>	<u>48 weeks</u>	<u>24 weeks</u>	<u>48 weeks</u>
HCV-Genotype 1 and $\leq 2$ million copies/mL	32%	33%	4%	25%
HCV-Genotype 1 and $> 2$ million copies/mL	10%	27%	1%	33%
All HCV-Genotypes 1	17%	29%	2%	9%

10 The sustained virologic response for all HCV genotype 1 patients treated for 24 and 48 weeks with the Combination Therapy was statistically significantly superior to that observed for all HCV genotype 1 patients treated with INTRON A plus placebo for 24 and 48 weeks. The sustained virologic response for HCV-genotype 1 patients with a baseline HCV viral load of greater than 2 million copies/mL was statistically significantly superior at 24 and 48 weeks of Combination Therapy compared to INTRON A + Placebo for 24 and 48 weeks.

**Table 17. Sustained Virologic Response by HCV -  
Genotype and Baseline HCV-RNA Levels for Studies 1 & 2**

<u>Patients</u>	<u>Intron A + Ribavirin</u>		<u>Intron A + Placebo</u>	
	<u>24 weeks</u>	<u>48 weeks</u>	<u>24 weeks</u>	<u>48 weeks</u>
HCV Genotype 1 and Baseline HCV-RNA levels of $\leq 2$ million copies/mL	32%	33%	4%	25%
HCV Genotype 1 and Baseline HCV-RNA levels of $> 2$ million copies/mL	11%	27%	1%	4%
Other HCV Genotypes <sup>1</sup> and Baseline HCV-RNA levels of $> 2$ million	61%	64%	25%	36%
Other HCV Genotypes <sup>1</sup> and Baseline HCV-RNA levels of $> 2$ million copies	62%	61%	11%	10%

5

<sup>1</sup> HCV-genotypes 4/5/6 account for 3% of patients Sustained Response for I+R :  
24% (24 wks) and 38% (48 weeks) for I+P : 0% (24 weeks) and 11% (48 weeks)  
Sustained Response Rate for HCV-Genotypes 2+3 is same as above

10

**Table 18. Sustained Virologic Response by  
Baseline HCV-RNA Levels for ALL Patients (Studies 1 & 2)**

<u>Patients</u>	<u>Intron A + Ribavirin</u>		<u>Intron A + Placebo</u>	
	<u>24 weeks</u>	<u>48 weeks</u>	<u>24 weeks</u>	<u>48 weeks</u>
Patients with Baseline HCV-RNA Levels $\leq 2$ million copies/mL	44%	46%	9%	36%
Patients with Baseline HCV-RNA Levels $> 2$ million copies/mL	27%	38%	4%	10%

**Table 19. Percent of Sustained Virologic Responders by  
Time to First No Detectable HCV-RNA Levels for Studies 1 & 2**

5

<u>Time to First No-Detectable HCV-RNA (Weeks)</u>	<u>Intron A + Ribavirin</u>		<u>Intron A + Placebo</u>	
	<u>24 weeks</u>	<u>48 weeks</u>	<u>24 weeks</u>	<u>48 weeks</u>
4	83%(92/111)	82%(94/115)	48%(10/21)	71%(47/66)
12	44%(66/149)	66%(91/137)	9%(3/32)	35%(29/84)
24	19%(8/42)	44%(20/45)	0%(0/22)	15%(6/39)

**Table 20. ALT Response: Normalization of Serum ALT in All Patients from Studies 1 & 2**

	INTRON A + Ribavirin		INTRON A + Placebo			
At End of Treat- ment	A 24 weeks (N=505)	B 48 weeks (N=505)	C 24 weeks (N=231)	D 48 weeks (N=503)	P <sup>a</sup> Value B vs D	P <sup>a</sup> Value A vs D
<b><u>Study</u></b>						
1 & 2 <sup>b</sup>	66% (329)	66% (334)	24% (56)	37% (185)	<0.001	0.739
1	58% (133)	61% (138)	24% (56)	28% (62)	<0.001	<0.001
2	71% (196)	71% (196)	–	44% (123)	<0.001	<0.001
<b><u>End of Follow- Up</u></b>						
<b><u>Study</u></b>						
1 + 2 <sup>b</sup>	36% (181)	44% (221)	11% (25)	24% (102)	<0.001	<0.001
1	32% (72)	36% (83)	11% (25)	16% (35)	<0.001	<0.001
2	39% (109)	50% (138)	–	24% (67)	<0.001	<0.001

5

<sup>a</sup> Cochran-Mantel-Haenszel general association for combined results; Fisher's Exact Test for individual studies.

<sup>b</sup> Combined results

10

Many modifications and variations of this invention can be made without departing from its spirit and scope, as will be apparent to those skilled in the art. The specific embodiments described herein are offered

5 by way of example only, and the invention is to be limited only by the terms of the appended claims, along with the full scope of equivalents to which such claims are entitled.

### CLAIMS

1. The use of ribavirin for the manufacture of a pharmaceutical  
5 composition for treating an antiviral treatment naive patient having  
chronic hepatitis C infection to eradicate detectable HCV-RNA by a  
method comprising administering an effective amount of ribavirin in  
association with an effective amount of interferon alpha for a time period  
of 20 up to 50 weeks; and wherein the antiviral treatment naive patient  
10 has a HCV genotype 2 or 3 infection, the administration of ribavirin in  
association with interferon alpha is effected for a time period of 20 - 30  
weeks, preferably 24 weeks. and wherein the antiviral treatment naive  
patient has a HCV genotype 1 infection administration of ribavirin in  
association with interferon alpha is effected for a time period of 40 - 50  
15 weeks preferably 48 weeks.

2. The use of interferon alpha for the manufacture of a pharmaceutical  
composition for treating an antiviral treatment naive patient having  
chronic hepatitis C infection to eradicate detectable HCV-RNA by a  
20 method comprising administering an effective amount of alpha interferon  
alpha in association with an effective amount of ribavirin for a time period  
of 20 to 50 weeks; and wherein the antiviral treatment naive patient has a  
HCV genotype 1 infection, administration of ribavirin in association with  
interferon alpha is effected for a time period of 40 - 50 weeks, preferably  
25 48 weeks and wherein the antiviral treatment naive patient has a HCV  
genotype 2 or 3 infection, the administration of ribavirin in association  
with interferon alpha is effected for a time period of 20 - 30 weeks,  
preferably 24 weeks.

3. The use of both ribavirin and alpha interferon alpha for the manufacture of pharmaceutical compositions for treating an antiviral treatment naive patient having chronic hepatitis C infection, to eradicate detectable HCV-RNA by a method comprising administering an effective amount of ribavirin in association with an effective amount of alpha-interferon for a time period of 20 up to 50 weeks; and wherein the antiviral treatment naive patient has a HCV genotype 1 infection administration of ribavirin in association with interferon alpha is effected for a time period of 40 - 50 weeks, preferably 48 weeks; and wherein the antiviral treatment naive patient has a HCV genotype 2 or 3 infection, the administration of ribavirin in association with interferon alpha is effected for a time period of 20 - 30 weeks, preferably 24 weeks.

4. The use as claimed in any preceding claim wherein the antiviral treatment naive patient is one having a HCV genotype 1 infection and a viral load of greater than 2 million copies per ml of HCV-RNA quantitative PCR.

5. The use in any one of claims 1, 2 or 3 wherein the patient has a HCV genotype 1 infection and a viral load of greater than 2 million copies per mL of HCV-1 RNA as measured by HCV-RNA quantitative PCR and wherein administration is effected for a time period of 40 - 50 weeks, preferably 48 weeks.

6. The use of both ribavirin and interferon alpha for the manufacture of pharmaceutical compositions for treating an antiviral treatment naive patient having chronic hepatitis C genotype 1 infection to eradicate detectable HCV-1-RNA by a method comprising administering an effective amount of ribavirin in association with an effective amount of interferon alpha for a time period of 40 - 50 weeks, preferably 48 weeks.



7. The use of both ribavirin and alpha interferon for the manufacture of pharmaceutical compositions for treating an antiviral treatment naive patient having chronic hepatitis C genotype 2 or 3 infection to eradicate detectable HCV-RNA by a method comprising administering an effective amount of ribavirin in association with an effective amount of alpha-interferon for a time period of 20 - 30 weeks, preferably 24 weeks.

8. The use of both ribavirin and alpha interferon for the manufacture of pharmaceutical compositions for treating an antiviral treatment naive patient having chronic hepatitis C genotype 1 infection, and a viral load of greater than 2 million copies per mL of HCV-1 RNA as measured by HCV-RNA quantitative PCR to eradicate detectable HCV-RNA by a method comprising administering an effective amount of ribavirin in association with an effective amount of alpha-interferon HCV for a time period of 40 - 50 weeks preferably 48 weeks.

9. The use as claimed in any one of the preceding claims wherein the amount of ribavirin administered is 800 to 1200 mg per day.

10. The use as claimed in any one of claims 1 to 9 wherein the interferon-alpha administered is selected from interferon alpha-2a, interferon alpha-2b, a consensus interferon, a purified interferon alpha product, a pegylated interferon alpha-2a or a pegylated interferon alpha-2b.

11. The use as claimed in any one of claims 1 to 9 wherein the alpha-interferon employed is an alpha-2b interferon and the amount of alpha-interferon administered is from 2 to 10 million IU per week on a weekly, TIW, QOD or daily basis.

12. The use as claimed in any one of claims 1 to 9 wherein the alpha interferon is alpha-2a interferon and the amount of alpha interferon administered is from 2 to 10 million IU per week on a weekly, TIW, QOD or daily basis.

# INTERNATIONAL SEARCH REPORT

National Application No  
PCT/US 99/07037

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 6 A61K38/21 A61K31/70 //(A61K38/21,31:70)

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	L CHEMELLO ET AL: "RESPONSE TO RIBAVIRIN, TO INTERFERON AND TO A COMBINATION OF BOTH IN PATIENTS WITH CHRONIC HEPATITIS C AND ITS RELATION TO HCV GENOTYPES" JOURNAL OF HEPATOLOGY, vol. 21, no. SUPPL. 01, 1 January 1994 (1994-01-01), page S12 XP002094476 the whole document ----- -/--	1-3,7, 9-12

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

27 July 1999

Date of mailing of the international search report

09/08/1999

Name and mailing address of the ISA  
European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Seegert, K

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 99/07037

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>BRACONIER J H ET AL: "COMBINED ALPHA-INTERFERON AND RIBAVIRIN TREATMENT IN CHRONIC HEPATITIS C: A PILOT STUDY" SCANDINAVIAN JOURNAL OF INFECTIOUS DISEASES, vol. 27, no. 4, 1 January 1995 (1995-01-01), pages 325-329, XP000600916 ISSN: 0036-5548 abstract page 327; table I</p> <p>---</p>	1-3,7, 9-12
X	<p>REICHARD O. ET AL: "Randomised, double-blind, placebo-controlled trial of interferon alpha-2b with and without ribavirin for chronic hepatitis C" THE LANCET, vol. 351, no. 9096, 10 January 1998 (1998-01-10), pages 83-87, XP002110247 cited in the application page 83 page 84; table 1</p> <p>---</p>	1-3,7, 9-12
P,X	<p>EP 0 903 148 A (SCHERING CORP) 24 March 1999 (1999-03-24) page 2 - page 3</p> <p>---</p>	1-3,7, 9-12
Y	<p>EP 0 707 855 A (SCHERING CORP) 24 April 1996 (1996-04-24) claims</p> <p>---</p>	1-12
Y	<p>BIZOLLON T. ET AL: "Pilot study of the combination of interferon alfa and ribavirin as therapy of recurrent hepatitis C after liver transplantation" HEPATOLOGY, vol. 26, no. 2, 1997, pages 500-504, XP002110248 abstract "Discussion" page 503</p> <p>---</p>	1-12
P,X	<p>MCHUTCHISON J.G. ET AL: "Interferon alfa-2b alone or in combination with ribavirin as initial treatment for chronic hepatitis C" NEW ENGLAND JOURNAL OF MEDICINE, vol. 339, no. 21, 19 November 1998 (1998-11-19), pages 185-1492, XP002110249 abstract</p> <p>---</p> <p style="text-align: center;">-/--</p>	1-12

# INTERNATIONAL SEARCH REPORT

national Application No  
PCT/US 99/07037

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>POL S. ET AL: "Ribavirin + Interferon-alfa 2B combination shows increased efficiency compared to high doses alfa INF, alone in naive patients infected by genotype 1B" HEPATOLOGY, vol. 28, no. 4 Part 2, October 1998 (1998-10), page 507a XP002110250 1378 abstract</p>	1-12
P,X	<p>POYNARD T. ET AL: "A International randomized trial of interferon alfa-2B and ribavirin (Intron A/Rebetol) 48 or 24 weeks vs Intron A 48 weeks for first line treatment of chronic hepatitis C" HEPATOLOGY, vol. 28, no. 4 Part 2, October 1998 (1998-10), page 387 XP002110251 900 abstract</p>	1-12

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/07037

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0903148 A	24-03-1999	AU 9473798 A WO 9915194 A	12-04-1999 01-04-1999
EP 0707855 A	24-04-1996	AU 5919296 A CA 2221314 A CN 1190895 A CZ 9703654 A HU 9802324 A JP 10506640 T NO 975309 A PL 323477 A SK 155997 A WO 9636351 A	29-11-1996 21-11-1996 19-08-1998 15-07-1998 28-05-1999 30-06-1998 19-11-1997 30-03-1998 07-10-1998 21-11-1996